

LOWER UMPQUA HOSPITAL OUTPATIENT NURSING DEPARTMENT 600 Ranch Road, Reedsport, OR 97467

For Referrals: (541) 271-2163 – Press #3 (all other services) then press #2 (Referrals)

For Outpatient Nursing: (541) 271-2171 ext. 5205

FAX: (541) 271-5433

REMDESIVIR THERAPY ORDERS (ADULT)

Patient Name		DOB
Patient Phone #	Patient Weight (Kg)) Patient Height (cm):
Patient Allergies		
		NPI#
ICD-10 Code (REQUIRED) _	J (Code
Primary Diagnosis		
Duration (or # of treatments	s): Ant	ticipated Infusion Date
 Lab orders should NOT responsibility of the or 	ce authorization has been initiated. be included on this form – place orders	s via usual method. Lab monitoring is the will need to be placed.
ORDERING GUIDELINES: ☑ Send FACE SHEET a	nd H&P or most recent chart note	
•	ry SARS-CoV-2 Infection ent with high risk of progression to seve	ere illness
MEDICATIONS:		
Loading Dose: ☑ Remdesivir 200 MG I	VPB on DAY 1 (100 mg / 0.9% sodiun	m chloride 100 mL at 100 mL/hour x 2 doses)
Scheduled Dosing: ☑ Remdesivir (100 MG	/ 0.9% sodium chloride 100 mL) 100	mg IVPB at 100 mL/hour DAILY on DAY 2 and DAY 3
clotting (alteplase), a ☑ alteplase (CATHFLO ☑ heparin, porcine Pres ☑ 0.9% sodium chlorid	and procedures for all vascular access nd/or dressing changes. <i>ACTIVASE</i>) injection 2 MG/2 mL, intra- servative Free (PF) 100 units/mL IV syr e 10 mL IV PRN flush as needed.	ringe, 500 units, intra-catheter x 1 PRN line care.
⋈ If applicable, may rem	ove PICC line at the completion of cou	urse of therapy. Continued next page →
Date Tim	ne Provider Signature _	
		patients to Lower Umpqua Hospital Outpatient Nursing Department, rs. as well as administration of any 340B drugs, remains with Lower

Page | 1 of 2 pages

Umpqua Hospital."



LOWER UMPQUA HOSPITAL OUTPATIENT NURSING DEPARTMENT 600 Ranch Road, Reedsport, OR 97467

For Referrals: (541) 271-2163 – Press #3 (all other services) then press #2 (Referrals)

For Outpatient Nursing: (541) 271-2171 ext. 5205

FAX: (541) 271-5433

REMDESIVIR THERAPY ORDERS (ADULT)

Patier	nt Name DOB
EMER	GENCY MEDICATIONS FOR HYPERSENSITIVITY / INFUSION REACTION:
	If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately 0.9% sodium chloride 200 mL/hour IV continuous infusion PRN hypersensitivity/infusion reaction
	acetaminophen (TYLENOL) tablet 650 MG, PO, Q4HRS PRN hypersensitivity/infusion reaction
X	Oxygen Therapy, low-flow, nasal cannula at 2 Liters/minute, may wean flow to discontinuation. SpO2 Goal – greater than/equal to 92%, PRN hypersensitivity/infusion reaction
X	diphenhydramine (BENADRYL) injection 25 MG IV x 1 PRN hypersensitivity/infusion reaction {Maximum of 3 doses}
	famotidine (PEPCID) 20 MG IV push over at least 2 minutes x 1 PRN hypersensitivity/infusion reaction
	hydrocortisone (SOLU-Medrol) 125 MG IV x 1 PRN hypersensitivity/infusion reaction
	albuterol (<i>PROVENTIL</i>) nebulizer solution 2.5 MG/3 mL, nebulization x 1 PRN wheezing, hypersensitivity/infusior reaction
X	epinephrine (ADRENOLIN) injection 0.3 MG/0.3 mL IM x 1 PRN anaphylaxis, hypersensitivity/ infusion reaction

"Statement of Responsibility of Parties: referring Prescriber agrees that in referring patients to Lower Umpqua Hospital Outpatient Nursing Department, the responsibility for the care related to these Outpatient Nursing Therapy Plan orders, as well as administration of any 340B drugs, remains with Lower Umpqua Hospital."

Date _____ Time _____ Provider Signature _____